

CHARGE FLUCTUATION IN HEAVY ION COLLISIONS *

FRITZ W. BOPP[†] AND JOHANNES RANFT

Universität Siegen, Fachbereich Physik, D-57068 Siegen, Germany

Charge fluctuations observed in early fixed-target proton-proton experiments are consistent with string models. In central heavy ion events the picture can change in two ways: strings can interact and find new ways to hadronize or they can be effectively inactivated to lose their dynamical role as ordering mechanism. Widely different charge fluctuations can be expected. The dispersion of the charges in a central rapidity box is an advantageous measure. In an explicit Dual-Parton-Model calculation using the DPMJET code and a randomized modification to simulated charge equilibrium, various energies and different nuclear sizes were considered. Local fluctuations were found to be a serious problem. However, for large enough detection regions charged particle fluctuations can provide a clear signal reflecting the basic dynamics of central heavy ion processes.

Charge Fluctuations in Fixed Target Hadron-Hadron Experiments

At fixed target experiments in hadronic multi-particle production it was possible to measure all charges of forward particles. In this way significant results could be obtained even with low energies available at the seventies¹:

- The charge fluctuations involve a restricted rapidity range.
- Qualitative agreement was obtained the Quigg-Thomas relation.

The Quigg-Thomas relation² was initially based on intermediately produced neutral clusters^{3,4}. It postulates for charge fluctuation across a rapidity y boundary

$$\langle \delta Q_{>y}^2 \rangle = \langle (Q_{>y} - \langle Q_{>y} \rangle)^2 \rangle = c \cdot dN_{\text{charge}}^{\text{non leading}}/dy. \quad (1)$$

To quantitatively fit the constant c with known resonances links with q resp. $\bar{q}$ exchanges had to be added⁵.

Such links appear in string models. We re-checked this old result using the Dual Parton Model code DPMJET⁶: For pp-scattering at laboratory energies of 205 GeV good agreement is obtained.

*Talk given at the Symposium on Multiparticle Dynamics, Alushta, Ukraine, Sept. 2002

[†]Partial support from the intas grant 97-31696 is acknowledged.

Strings as Ordering Mechanism

In QED real and virtual soft or collinear emissions cancel as the final states cannot be distinguished in measurement. For QCD such contributions involve long time scales leaving the perturbative regime. With the comparatively compact hadronic final states the emissions now cancel as the final states are equal.

In string models the hadronic final state is thought to be composed of color singlets called strings. If different soft or collinear contributions to a string production amplitude are summed their phases can lead to cancellations. In this way strings can act as infrared regulators.

It is possible that the usual soft phenomenology would emerge as extension of PQCD, if these cutoffs could be properly implemented. Strings can play an essential role as ordering mechanism of the dynamics of their production.

What changes for Heavy Ion Scattering?

With more interactions per nucleon strings will get more numerous and shorter. There are two possible quite distinct consequences:

- Denser *strings should interact* and find a different, possibly more efficient way to hadronize.
- A very large number of interactions can be expected to essentially *destroy the strings* as infrared regulator or ordering mechanism. The ensemble needed to describe the scattering then involves a much larger number of states.

Reasonable expectations for both cases are respectively:

- a *reduction* in density, an increase in baryon pairs and in strangeness⁷,
- an *increase* in density, possibly looking like local thermalization.

RHIC data seem to favor the first option. Unfortunately explicit models show large uncertainties. Clarification can come from charge fluctuation measurements.

Charge Fluctuations in Heavy-Ion Scattering Experiments

In heavy ion experiments the charge distribution of the particle contained in a central box with a given rapidity range $[-y_{\max.}, +y_{\max.}]$ can be measured and the dispersion of this distribution $\langle \delta Q^2 \rangle$ can be obtained to sufficient accuracy. In comparison to the fluctuations in the forward backward charge distributions the charge distribution into a central box (having to have two sufficiently separated borders) can be expected to require roughly twice the rapidity range to obtain information about long range charge flow.

Within the framework of equilibrium models the dispersion was proposed to distinguish between particles emerging from an equilibrium quark-gluon gas or from an equilibrium hadron gas^{8,9}. It should be pointed out that this estimate is not without theoretical problems^{10,11} having to do with the hadronisation process.

Besides the classic charge dispersion

$$\langle \delta Q^2 \rangle = \langle (Q - \langle Q \rangle)^2 \rangle \quad (2)$$

where $Q = N_+ - N_-$ is the net charge inside the box, it was proposed to just measure the mean standard deviation of the ratio R of positive to negative particles or the ratio F of the net charge to the total number of charged particles in the box. The motivation for choosing these ratios was to reduce the dependence of multiplicity fluctuations caused by the event structure. These quantities have problems for hadron-hadron or non-central heavy-ion events^{11,13}. As any conclusion will have to depend on a comparison of central processes with minimum bias and proton-proton events, there is a clear advantage to stick to the dispersion of the net charge distribution.

The ϕ - and Γ - measures also considered¹⁴ are closely related to $\langle \delta Q^2 \rangle$.

Quark Line Structure and Fluctuations in the Charge Flow

To visualize the meaning of charge flow measurements it is helpful to introduce the hypothesis that the flavor distribution of individual quarks factorizes. It is an adequate approximation, especially if long range fluctuations are considered.

The hypothesis leads to a *generalization of the Quigg-Thomas relation*⁵ determining the correlation of the charges exchanged across two arbitrary boundaries. A combination of such correlations yields the fluctuation of the charge within a $[-y_{\max.}, +y_{\max.}]$ box:

$$\langle \delta Q[\text{box}]^2 \rangle = n_{\text{entering box}}^{\text{quark lines}} \langle (\delta q)^2 \rangle \quad (3)$$

where $Q[\text{box}]$ is the charge in the box, where $n_{\text{entering box}}^{\text{quark lines}}$ is the number of quark lines entering or leaving the box, and where q is the charge of the quark on such a line. With the notation: $\delta Q = Q - \langle Q \rangle$, values $\langle \delta q^2 \rangle = 0.22 \dots 0.25$ can be obtained.

The relation allows to easily evaluate simple situations like the *thermalized limit* of a small box with an infinite reservoir outside. In an “*hadron gas case*” all particles contain two independent quarks coming from outside; in the so-called “*quark gluon gas case*” one of the quarks of each meson comes from the outside the other is ignored as a local hadronisation affair (one ignores $\langle q \rangle \neq 0$).

Expanding Box

For a *tiny box* — considering only at the first order in Δy — one trivially obtains the hadron gas value $\langle \delta Q^2 \rangle / \langle N_{\text{charged}} \rangle = 1$. If the box size increases to *one or two units of rapidity* on each side this ratio will decrease, as realistic models contain a short range component in the charge fluctuations.

After a box size passed the short range the *decisive region* starts. In all global equilibrium models the ratio will have to reach a flat value.

If a *large box* involves a significant part of the phase space the overall charge conservation has to be considered with a correction factor $\propto 1 - y_{\max.}/Y_{\text{kin.max.}}$. At present energies the decisive and large region are not separated.

String Model Predictions

Charges are locally compensated as the range spanned by quark lines in links or during resonance decays is limited. The total contribution will be determined by the density of quark lines reflecting the *number of strings* at the boundaries:

$$\langle \delta Q^2 \rangle \propto \rho_{\text{charged}}(y_{\text{max}}). \quad (4)$$

This resulting scaling is illustrated in a DPMJET⁶ comparison between both quantities in (4) shown in Fig. 1 for RHIC and LHC energies. For smaller boxes




Figure 1. The dispersion of the charge distribution and the density on the box boundary for central gold gold resp. lead lead scattering at RHIC and LHC energies.

there is a correction as some of the quark lines intersect both boundaries. For large rapidity sizes there is a minor increase from the leading charge flow Q_L originating in the incoming particles¹². In a more careful consideration³ one can subtract this contribution $\langle Q_L \rangle (1 - \langle Q_L \rangle)$ and concentrate truly on the fluctuation. A simple estimate — with a width of neighboring string break ups and a width from resonance decays — leads to consistent values¹¹.

A comparable result was obtained for the proton-proton case¹¹.

String Model versus “Hadron Gas”

It was argued⁹ that the experimental results should be “purified” to account for charge conservation. We prefer a reference model with a posteriori *randomized charges*. This unbiased method can be obviously also directly applied to experimental data. Using DPMJET for RHIC and LHC energies for *proton-proton* and *central lead-lead* collisions we obtain the “statistical” prediction shown in Fig. 2.

We employed the correction factor $(1 - \int_0^{y_{\text{max}}} \rho_{\text{charge}} dy) / \int_0^{Y_{\text{kin.max}}} \rho_{\text{charge}} dy$ proposed by⁹ to check consistency and obtained expected the flat distribution.




Figure 2. Charge fluctuations with a posteriori randomized charges for p-p scattering and the most central 5% in Pb-Pb scattering at RHIC energies ($\sqrt{s} = 200$ A GeV) and at LHC energies ($\sqrt{s} = 6000$ A GeV). The results are also shown with a correction factor to account for the overall charge conservation.




Figure 3. Comparison of the charge fluctuations obtained in a string model DP-MJET with a model using a posteriori randomized charges for p-p scattering and the most central 5% in Pb-Pb scattering at RHIC energies ($\sqrt{s} = 200$ A GeV) and at LHC energies ($\sqrt{s} = 6000$ A GeV).

Taking the DPMJET string model and the randomized “hadron gas” version as extreme cases the decisive power can be tested. As shown in Fig.3 there is a *measurable* distinction at RHIC energies and *sizable* one at LHC energies.

The spectra change roughly by a factor of 400 between simple proton-proton scattering and central lead lead scattering. The surprising similarity between p-p and Pb-Pb in the Fig. 3 can be understood as collective effects to a large part not included in the model. Also no dependence on the centrality was observed in DPMJET for Pb-Pb scattering at RHIC energies ($\sqrt{s} = 200$ A GeV)¹¹. This experimentally measurable centrality dependence allows to directly observe collective effects without reference to a particular model.

Conclusion

The dispersion of the charge distribution in a central box of varying size is an extremely powerful measure. It allows to directly and quantitatively test *the presence of equilibrizing processes* and remaining *dynamical corrections* to equilibrated distributions.

References

1. U. Idschok et al. [Bonn-Hamburg-Munich Collaboration], Nucl. Phys. **B67**, 93 (1973), A. Bialas, K. Fialkowski, M. Jezabek and M. Zielinski, Acta Phys. Polon. B **6**, 39 (1975), J. Whitmore, Phys. Rept. **27**, 187 (1976), L. Foa, Phys. Rept. **22**, 1 (1975), R. Brandelik *et al.* [TASSO Collaboration], Phys. Lett. B **100**, 357 (1981), C. Quigg, Phys. Rev. **D12**, 834 (1975).
2. C. Quigg and G. H. Thomas, Phys. Rev. **D7**, 2757 (1973), C. Quigg, Phys. Rev. **D12**, 834 (1975).
3. F. W. Bopp, Riv. Nuovo Cim. **1**, 1 (1978).
4. J. Ranft, Fortsch. Phys. **23**, 467 (1975), S. P. Misra and B. K. Parida, Pramana **20**, 375 (1983), S. P. Misra and B. K. Parida, Pramana **20**, 375 (1983), T. T. Chou and C. N. Yang, Phys. Rev. **D7**, 1425 (1973), J. L. Newmeyer and D. Sivers, Phys. Rev. D **10**, 204 (1974), K. F. Loe, K. K. Phua and S. C. Chan, Lett. Nuovo Cim. **18**, 137 (1977), E. N. Argyres and C. S. Lam, Phys. Rev. D **16**, 114 (1977), C. B. Chiu and K. Wang, Phys. Rev. D **13**, 3045 (1976), J. Dias de Deus and S. Jadach, Phys. Lett. B **66**, 81 (1977), M. Jezabek, Phys. Lett. B **67**, 292 (1977).
5. R. Baier and F. W. Bopp, Nucl. Phys. **B79**, 344 (1974), P. Aurenche and F. W. Bopp, Nucl. Phys. **B119**, 157 (1977).
6. J. Ranft, Phys. Rev. **D 510** 64 (1995); J. Ranft, hep-ph/9911213 (Siegen preprint SI-99-5); J. Ranft, hep-ph/9911232 (Siegen preprint SI-99-6).
7. N. S. Amelin, N. Armesto, C. Pajares and D. Sousa, Eur. Phys. J. C **22**, 149 (2001) [arXiv:hep-ph/0103060].
8. M. Asakawa, U. Heinz and B. Muller, Phys. Rev. Lett. **85**, 2072 (2000), S. Jeon and V. Koch, Phys. Rev. Lett. **85**, 2076 (2000) [hep-ph/0003168], S. Jeon and V. Koch, Phys. Rev. Lett. **83**, 5435 (1999) [nucl-th/9906074].
9. M. Bleicher, S. Jeon and V. Koch, Phys. Rev. **C62**, 061902 (2000) [hep-ph/0006201]; V. Koch, M. Bleicher and S. Jeon, nucl-th/0103084, M. Bleicher *et al.*, J. Phys. **G25** (1999) 1859 [hep-ph/9909407].
10. F. W. Bopp, Nucl. Phys. **B191**, 75 (1981).
11. F. W. Bopp and J. Ranft, Acta Phys. Polon. B **33**, 1505 (2002) [arXiv:hep-ph/0204010]. F. W. Bopp and J. Ranft, Eur. Phys. J. C **22**, 171 (2001) arXiv:hep-ph/0105192, K. Fialkowski and R. Wit, hep-ph/0006023, "Charge fluctuations in a final state with QGP," hep-ph/0101258, A. Bialas, arXiv:hep-ph/0203047.
12. A. Capella, C. Merino and J. Tran Thanh Van, Z. Phys. C **43**, 663 (1989).
13. K. Adcox [PHENIX Collaboration], arXiv:nucl-ex/0203014.
14. S. Mrowczynski, Phys. Rev. C **66**, 024904 (2002), C. Pruneau, S. Gavin and S. Voloshin, Phys. Rev. C **66**, 044904 (2002) [arXiv:nucl-ex/0204011].